

THE MATHEMATICAL ASSOCIATION OF AMERICA
AMERICAN MATHEMATICS COMPETITIONS



24th Annual (*Alternate*)

AMERICAN INVITATIONAL
MATHEMATICS EXAMINATION
(AIME II)

Wednesday, March 22, 2006

1. DO NOT OPEN THIS BOOKLET UNTIL YOUR PROCTOR GIVES THE SIGNAL TO BEGIN.
2. This is a 15-question, 3-hour examination. All answers are integers ranging from 000 to 999, inclusive. Your score will be the number of correct answers; i.e., there is neither partial credit nor a penalty for wrong answers.
3. No aids other than scratch paper, graph paper, ruler, compass, and protractor are permitted. In particular, **calculators and computers are not permitted.**
4. A combination of the AIME and the American Mathematics Contest 10 or the American Mathematics Contest 12 scores are used to determine eligibility for participation in the U.S.A. Mathematical Olympiad (USAMO). The USAMO will be given on TUESDAY and WEDNESDAY, April 18 & 19, 2006.
5. Record all of your answers, and certain other information, on the AIME answer form. Only the answer form will be collected from you.

The publication, reproduction or communication of the problems or solutions of the AIME during the period when students are eligible to participate seriously jeopardizes the integrity of the results. Duplication at any time via copier, telephone, email, world wide web, or media of any type is a violation of the competition rules.


1. In convex hexagon $ABCDEF$, all six sides are congruent, $\angle A$ and $\angle D$ are right angles, and $\angle B$, $\angle C$, $\angle E$, and $\angle F$ are congruent. The area of the hexagonal region is $2116(\sqrt{2}+1)$. Find AB .
2. The lengths of the sides of a triangle with positive area are $\log_{10} 12$, $\log_{10} 75$, and $\log_{10} n$, where n is a positive integer. Find the number of possible values for n .
3. Let P be the product of the first 100 positive odd integers. Find the largest integer k such that P is divisible by 3^k .
4. Let $(a_1, a_2, a_3, \dots, a_{12})$ be a permutation of $(1, 2, 3, \dots, 12)$ for which

$$a_1 > a_2 > a_3 > a_4 > a_5 > a_6 \quad \text{and} \quad a_6 < a_7 < a_8 < a_9 < a_{10} < a_{11} < a_{12}.$$

An example of such a permutation is $(6, 5, 4, 3, 2, 1, 7, 8, 9, 10, 11, 12)$. Find the number of such permutations.

5. When rolling a certain unfair six-sided die with faces numbered 1, 2, 3, 4, 5, and 6, the probability of obtaining face F is greater than $1/6$, the probability of obtaining the face opposite face F is less than $1/6$, the probability of obtaining each of the other faces is $1/6$, and the sum of the numbers on each pair of opposite faces is 7. When two such dice are rolled, the probability of obtaining a sum of 7 is $47/288$. Given that the probability of obtaining face F is m/n , where m and n are relatively prime positive integers, find $m+n$.
6. Square $ABCD$ has sides of length 1. Points E and F are on \overline{BC} and \overline{CD} , respectively, so that $\triangle AEF$ is equilateral. A square with vertex B has sides that are parallel to those of $ABCD$ and a vertex on \overline{AE} . The length of a side of this smaller square is $\frac{a - \sqrt{b}}{c}$, where a , b , and c are positive integers and b is not divisible by the square of any prime. Find $a + b + c$.
7. Find the number of ordered pairs of positive integers (a, b) such that $a + b = 1000$ and neither a nor b has a zero digit.

8. There is an unlimited supply of congruent equilateral triangles made of colored paper. Each triangle is a solid color with the same color on both sides of the paper. A large equilateral triangle is constructed from four of these paper triangles as shown. Two large triangles are considered distinguishable if it is not possible to place one on the other, using translations, rotations, and/or reflections, so that their corresponding small triangles are of the same color. Given that there are six different colors of triangles from which to choose, how many distinguishable large equilateral triangles can be constructed?



9. Circles \mathcal{C}_1 , \mathcal{C}_2 , and \mathcal{C}_3 have their centers at $(0, 0)$, $(12, 0)$, and $(24, 0)$, and have radii 1, 2, and 4, respectively. Line t_1 is a common internal tangent to \mathcal{C}_1 and \mathcal{C}_2 and has a positive slope, and line t_2 is a common internal tangent to \mathcal{C}_2 and \mathcal{C}_3 and has a negative slope. Given that lines t_1 and t_2 intersect at (x, y) , and that $x = p - q\sqrt{r}$, where p , q , and r are positive integers and r is not divisible by the square of any prime, find $p + q + r$.
10. Seven teams play a soccer tournament in which each team plays every other team exactly once. No ties occur, each team has a 50% chance of winning each game it plays, and the outcomes of the games are independent. In each game, the winner is awarded 1 point and the loser gets 0 points. The total points are accumulated to decide the ranks of the teams. In the first game of the tournament, team A beats team B . The probability that team A finishes with more points than team B is m/n , where m and n are relatively prime positive integers. Find $m + n$.
11. A sequence is defined as follows: $a_1 = a_2 = a_3 = 1$, and, for all positive integers n , $a_{n+3} = a_{n+2} + a_{n+1} + a_n$. Given that $a_{28} = 6090307$, $a_{29} = 11201821$, and $a_{30} = 20603361$, find the remainder when $\sum_{k=1}^{28} a_k$ is divided by 1000.
12. Equilateral $\triangle ABC$ is inscribed in a circle of radius 2. Extend \overline{AB} through B to point D so that $AD = 13$, and extend \overline{AC} through C to point E so that $AE = 11$. Through D , draw a line ℓ_1 parallel to \overline{AE} , and through E , draw a line ℓ_2 parallel to \overline{AD} . Let F be the intersection of ℓ_1 and ℓ_2 . Let G be the point on the circle that is collinear with A and F and distinct from A . Given that the area of $\triangle CBG$ can be expressed in the form $p\sqrt{q}/r$, where p , q , and r are positive integers, p and r are relatively prime, and q is not divisible by the square of any prime, find $p + q + r$.
13. How many integers N less than 1000 can be written as the sum of j consecutive positive odd integers for exactly 5 values of $j \geq 1$?
14. Let S_n be the sum of the reciprocals of the nonzero digits of the integers from 1 to 10^n , inclusive. Find the smallest positive integer n for which S_n is an integer.
15. Given that x , y , and z are real numbers that satisfy

$$\begin{aligned} x &= \sqrt{y^2 - \frac{1}{16}} + \sqrt{z^2 - \frac{1}{16}} \\ y &= \sqrt{z^2 - \frac{1}{25}} + \sqrt{x^2 - \frac{1}{25}} \\ z &= \sqrt{x^2 - \frac{1}{36}} + \sqrt{y^2 - \frac{1}{36}}, \end{aligned}$$

and that $x + y + z = m/\sqrt{n}$, where m and n are positive integers, and n is not divisible by the square of any prime, find $m + n$.

Your Exam Manager will receive a copy of the 2006 AIME Solution Pamphlet with the scores.

CONTACT US -- Correspondence about the problems and solutions for this AIME and orders for any of our publications should be addressed to:

American Mathematics Competitions
University of Nebraska, P.O. Box 81606
Lincoln, NE 68501-1606

Phone: 402-472-2257; Fax: 402-472-6087; email: amcinfo@unl.edu

The problems and solutions for this AIME were prepared by the MAA's Committee on the AIME under the direction of:

David Hankin, AIME Chair
207 Corbin Place, Brooklyn, NY 11235 USA

2006 USAMO -- THE USA MATHEMATICAL OLYMPIAD (USAMO) is a 6-question, 9-hour, essay-type examination. The USAMO will be held in your school on Tuesday, April 18 & Wednesday, April 19. Your teacher has more details on who qualifies for the USAMO in the AMC 10/12 and AIME Teachers' Manuals. The best way to prepare for the USAMO is to study previous years of these exams, the World Olympiad Problems/Solutions and review the contents of the Arbelos. Copies may be ordered from the web sites indicated below.

PUBLICATIONS -- For a complete listing of available publications please visit the following web sites:

AMC - - <http://www.unl.edu/amc/d-publication/publication.html>

MAA -- https://enterprise.maa.org/ecomtpro/timssnet/common/tnt_frontpage.cfm

The American Mathematics Competitions

are Sponsored by

The Mathematical Association of America — MAA www.maa.org/
University of Nebraska – Lincoln — UN-L www.unl.edu/
The Akamai Foundation – www.akamai.com/

Contributors

American Mathematical Association of Two Year Colleges – AMATYC..... www.amatyc.org/
American Mathematical Society — AMS www.ams.org/
American Society of Pension Actuaries — ASPA..... www.aspa.org/
American Statistical Association — ASA..... www.amstat.org/
Art of Problem Solving — www.artofproblemsolving.com/
Canada/USA Mathcamp — C/USA MC www.mathcamp.org/
Canada/USA Mathpath — C/USA MP www.mathpath.org/
Casualty Actuarial Society — CAS www.casact.org/
Clay Mathematics Institute — CMI..... www.claymath.org/
Institute for Operations Research and the Management Sciences — INFORMS www.informs.org/
L. G. Balfour Company www.balfour.com/
Mu Alpha Theta — MAT www.mualphatheta.org/
National Assessment & Testing www.natassessment.com/
National Council of Teachers of Mathematics — NCTM..... www.nctm.org/
Pedagoguery Software Inc. — www.peda.com/
Pi Mu Epsilon — PME..... www.pme-math.org/
Society of Actuaries — SOA www.soa.org/
U. S. A. Math Talent Search — USAMTS www.usamts.org/
W. H. Freeman and Company www.whfreeman.com/
Wolfram Research Inc. www.wolfram.com/